

Coos County Community Development Attn: Planning 60 E. Second St. Coquille, OR 97423 August 11, 2023

Re: Supplemental Information for Incomplete Application ACU 23-034

Information requested in letter dated June 22, 2023, deeming the subject application incomplete:

(1) Plot Plan

Revised plot plan is attached to this document.

(2) List of involved parcels and acknowledgment of consolidation

Subject property: 28S-14W-20BB, Tax Lots 1103, 1104, 1100, 1101.

The proposed development is located primarily on TL 1104, with vehicular access through TL 1100 and 1101. No separate or additional development is proposed for the remaining parcels in this tract.

The property owner agrees to consolidate all four properties within the tract, and requests this be a condition of approval imposed on the request upon Conditional Use approval to develop the property as proposed.

(3) Soil report to address OAR 660-033-0020(8)(a) (4.6.200 #3C)

Site specific and vicinity NRCS Soil Web Surveys are attached to this document.

A percentage of the tract is composed of "8 B -Bullards Sandy Loam, 0-7 percent slope" soil which is classified as farmland with statewide significance. However, the areas of the tract with this soil classification are surrounded by "8 E – Bullards sandy loam, 30-50 percent slope" soil, which has no farmland significance and would be impractical for farm use due to the characteristics and properties of the soil, topography of the property, and the surrounding residential development.

Though the tract may have protection under OAR 660-033-0020 as significant agricultural land, the individual properties that comprise the subject tract would be impossible to farm due to their small one-acre sizes, lack of water rights, mixture of soils classifications, varying topographies, and assumed compatibility issues with operating a farm near established residential developments.

(4) Description of water rights (4.6.200 #3D)

The property owners have no knowledge or record of existing water rights to any portion of the subject property. The Oregon Water Resources Department (OWRD) has no record of previously granted water rights. Neither the individual lots, nor the tract, have water rights.

Additional:

The current and previous owners of record are lifelong residents of Coos County and own other property in close proximity that has been used for farming, specifically growing and harvesting cranberries, throughout the last 50 years. OWRD records reflect these properties and their history of farm use. The water rights granted to those properties do not extend or apply to the subject tract.

At one time, the subject tract was intended to be put into this farm use to support the family's farm operation; however, the issues raised in this application ultimately made any plan for farming the properties individually or as a tract, impossible.

Megan Lawrence

Owner, South Coast Consulting LLC

PROPOSED SITE PLAN FOR ZONING COMPLIANCE





USDA United States Department of Agriculture

> Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Coos County, Oregon

Kent



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	
Soil Map	9
Legend	10
Map Unit Legend	11
Map Unit Descriptions	11
Coos County, Oregon	13
8B—Bullards sandy loam, 0 to 7 percent slopes	13
8E—Bullards sandy loam, 30 to 50 percent slopes	14
Soil Information for All Uses	16
Suitabilities and Limitations for Use	16
Building Site Development	16
Dwellings Without Basements	16
Land Classifications	20
Farmland Classification	20
Soil Reports	26
Building Site Development	26
Dwellings and Small Commercial Buildings	
Land Classifications	28
Prime and other Important Farmlands	28
References	31

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



	MAP LEGEND			MAP INFORMATION
Area of Int	terest (AOI) Area of Interest (AOI)	8	Spoil Area Stony Spot	The soil surveys that comprise your AOI were mapped at 1:20,000.
Soils	Soil Map Unit Polygons Soil Map Unit Lines Soil Map Unit Points Point Features	00 0 0	Very Stony Spot Wet Spot Other Special Line Features	Warning: Soil Map may not be valid at this scale. Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed
() () () () () () () () () () () () () (Blowout Borrow Pit Clay Spot	Water Features Streams and Canals Transportation HHH Rails	tures Streams and Canals ation Rails	scale. Please rely on the bar scale on each map sheet for map measurements.
◇ ¥ *	Closed Depression Gravel Pit Gravelly Spot	}	Interstate Highways US Routes Major Roads	Source of Map: Natural Resources Conservation Service Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)
۵ بر ش	Landfill Lava Flow Marsh or swamp Mine or Quarry	Backgroui	Local Roads nd Aerial Photography	Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.
© 0 ~	Miscellaneous Water Perennial Water Rock Outcrop			This product is generated from the USDA-NRCS certified data as of the version date(s) listed below. Soil Survey Area: Coos County, Oregon
+ :: =	Saline Spot Sandy Spot Severely Eroded Spot			Survey Area Data: Version 18, Sep 8, 2022 Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.
۵ بو بو	Sinkhole Slide or Slip Sodic Spot			Date(s) aerial images were photographed: Oct 5, 2019—Oct 10, 2019 The orthophoto or other base map on which the soil lines were
				compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
8B	Bullards sandy loam, 0 to 7 percent slopes	2.5	56.0%
8E	Bullards sandy loam, 30 to 50 percent slopes	2.0	44.0%
Totals for Area of Interest		4.5	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however,

onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Coos County, Oregon

8B—Bullards sandy loam, 0 to 7 percent slopes

Map Unit Setting

National map unit symbol: 21rc Elevation: 30 to 600 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Bullards and similar soils: 75 percent *Minor components:* 9 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Bullards

Setting

Landform: Marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Parent material: Mixed eolian and marine deposits

Typical profile

Oi - 0 to 3 inches: slightly decomposed plant material

H1 - 3 to 10 inches: sandy loam

H2 - 10 to 44 inches: gravelly sandy loam

H3 - 44 to 63 inches: sand

Properties and qualities

Slope: 0 to 7 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: F004AC410OR - Coastal Upland Warm Forest Forage suitability group: Well Drained <15% Slopes (G004AY014OR) Other vegetative classification: Well Drained <15% Slopes (G004AY014OR) Hydric soil rating: No

Minor Components

Blacklock

Percent of map unit: 9 percent

Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: Yes

8E—Bullards sandy loam, 30 to 50 percent slopes

Map Unit Setting

National map unit symbol: 21rg Elevation: 50 to 600 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Not prime farmland

Map Unit Composition

Bullards and similar soils: 80 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Bullards

Setting

Landform: Marine terraces Landform position (three-dimensional): Riser Down-slope shape: Linear Across-slope shape: Linear Parent material: Mixed eolian and marine deposits

Typical profile

Oi - 0 to 3 inches: slightly decomposed plant material *H1 - 3 to 10 inches:* sandy loam *H2 - 10 to 44 inches:* gravelly sandy loam *H3 - 44 to 63 inches:* sand

Properties and qualities

Slope: 30 to 50 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6e Hydrologic Soil Group: B Ecological site: F004AC410OR - Coastal Upland Warm Forest Hydric soil rating: No

Soil Information for All Uses

Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

Building Site Development

Building site development interpretations are designed to be used as tools for evaluating soil suitability and identifying soil limitations for various construction purposes. As part of the interpretation process, the rating applies to each soil in its described condition and does not consider present land use. Example interpretations can include corrosion of concrete and steel, shallow excavations, dwellings with and without basements, small commercial buildings, local roads and streets, and lawns and landscaping.

Dwellings Without Basements

ENG - Engineering

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper.

The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification of the soil. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. "Not limited" indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. "Somewhat limited" indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. "Very limited" indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The map unit components listed for each map unit in the accompanying Summary by Map Unit table in Web Soil Survey or the Aggregation Report in Soil Data Viewer are determined by the aggregation method chosen. An aggregated rating class is shown for each map unit. The components listed for each map unit are only those that have the same rating class as listed for the map unit. The percent composition of each component in a particular map unit is presented to help the user better understand the percentage of each map unit that has the rating presented.

Other components with different ratings may be present in each map unit. The ratings for all components, regardless of the map unit aggregated rating, can be viewed by generating the equivalent report from the Soil Reports tab in Web Soil Survey or from the Soil Data Mart site. Onsite investigation may be needed to validate these interpretations and to confirm the identity of the soil on a given site.



	MAP LEGEND				MAP INFORMATION
Area of In	terest (AOI) Area of Interest (AOI)	Backgrou	nd Aerial Photography		The soil surveys that comprise your AOI were mapped at 1:20,000.
Soils					
Soil Rat	ing Polygons				Warning: Soil Map may not be valid at this scale.
	Very limited				Enlargement of maps beyond the scale of mapping can cause
	Somewhat limited				misunderstanding of the detail of mapping and accuracy of soil
	Not limited				line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed
	Not rated or not available				scale.
Soil Rat	ing Lines				
~	Very limited				Please rely on the bar scale on each map sheet for map
~	Somewhat limited				measurements.
~	Not limited				Source of Map: Natural Resources Conservation Service
	Not rated or not available				Coordinate System: Web Mercator (EPSG:3857)
Soil Rat	ing Points				
	Very limited				Maps from the Web Soil Survey are based on the Web Mercator
	Somewhat limited				distance and area. A projection that preserves area, such as the
	Not limited				Albers equal-area conic projection, should be used if more
	Not rated or not available				
	tures				This product is generated from the USDA-NRCS certified data as
~	Streams and Canals				of the version date(s) listed below.
Transport	ation				Soil Survey Area: Coos County, Oregon
+++	Rails				Survey Area Data: Version 18, Sep 8, 2022
~	Interstate Highways				Soil map units are labeled (as space allows) for map scales
~	US Routes				1:50,000 or larger.
~	Major Roads				Date(s) aerial images were photographed: Oct 5, 2019—Oct 10
~	Local Roads				2019
					The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Tables—Dwellings V	Without Basements
--------------------	--------------------------

Map unit symbol	Map unit name	Rating	Component name (percent)	Rating reasons (numeric values)	Acres in AOI	Percent of AOI
8B	Bullards sandy loam, 0 to 7 percent slopes	Not limited	Bullards (75%)		2.5	56.0%
8E	Bullards sandy loam, 30 to 50 percent slopes	Very limited	Bullards (80%)	Slope (1.00)	2.0	44.0%
Totals for Area of	f Interest				4.5	100.0%

Rating	Acres in AOI	Percent of AOI
Not limited	2.5	56.0%
Very limited	2.0	44.0%
Totals for Area of Interest	4.5	100.0%

Rating Options—Dwellings Without Basements

Aggregation Method: Dominant Condition Component Percent Cutoff: None Specified Tie-break Rule: Higher

Land Classifications

Land Classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Farmland Classification

Farmland classification identifies map units as prime farmland, farmland of statewide importance, farmland of local importance, or unique farmland. It identifies the location and extent of the soils that are best suited to food, feed, fiber, forage, and oilseed crops. NRCS policy and procedures on prime and unique farmlands are published in the "Federal Register," Vol. 43, No. 21, January 31, 1978.





Custom Soil Resource Report

Prime farmland if Farmland of statewide Farmland of statewide Farmland of unique Prime farmland if 1 A الريادي -----subsoiled, completely importance, if drained and importance, if irrigated importance subsoiled, completely removing the root either protected from and reclaimed of excess removing the root Not rated or not available $\mathcal{F}^{(1)}(\mathcal{F})$ inhibiting soil layer flooding or not frequently salts and sodium inhibiting soil layer flooded during the Soil Rating Points Prime farmland if irrigated Farmland of statewide Prime farmland if arowing season and the product of I (soil importance, if drained or irrigated and the product Not prime farmland erodibility) x C (climate Farmland of statewide either protected from of I (soil erodibility) x C factor) does not exceed importance, if irrigated flooding or not frequently All areas are prime (climate factor) does not and drained flooded during the farmland exceed 60 60 growing season Prime farmland if irrigated Farmland of statewide Prime farmland if drained Prime farmland if -الجريداتين and reclaimed of excess importance, if irrigated Farmland of statewide irrigated and reclaimed -Prime farmland if salts and sodium and either protected from importance, if warm of excess salts and protected from flooding or flooding or not frequently enough, and either sodium Farmland of statewide ----not frequently flooded flooded during the drained or either Farmland of statewide importance during the growing growing season protected from flooding or importance Farmland of statewide not frequently flooded season a 🖬 Farmland of statewide Farmland of statewide importance, if drained during the growing Prime farmland if irrigated importance, if subsoiled. importance, if drained Farmland of statewide season completely removing the importance, if protected Prime farmland if drained Farmland of statewide root inhibiting soil layer Farmland of statewide from flooding or not and either protected from importance, if protected importance, if warm Farmland of statewide 100 frequently flooded during flooding or not frequently from flooding or not enough importance, if irrigated the growing season flooded during the frequently flooded during and the product of I (soil Farmland of statewide growing season the growing season Farmland of statewide 1990 B erodibility) x C (climate importance, if thawed importance, if irrigated Prime farmland if irrigated Farmland of statewide factor) does not exceed Farmland of local 1000 and drained importance, if irrigated 60 importance Prime farmland if irrigated Farmland of local ----and either protected from importance, if irrigated flooding or not frequently flooded during the growing season

Custom Soil Resource Report

Farmland of statewide importance, if drained and either protected from flooding or not frequently		Farmland of statewide importance, if irrigated and reclaimed of excess salts and sodium		Farmland of unique importance Not rated or not available	The soil surveys that comprise your AOI were mapped at 1:20,000.
flooded during the growing season		Farmland of statewide	Water Feat	tures Streams and Canals	Warning: Soil Map may not be valid at this scale.
Farmland of statewide importance, if irrigated		either protected from flooding or not frequently	Transporta	ation	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil
and drained Farmland of statewide	_	growing season	••••	Rails	line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed
and either protected from flooding or not frequently		importance, if warm enough, and either	~	US Routes	scale.
flooded during the growing season		drained or either protected from flooding or	~	Major Roads	Please rely on the bar scale on each map sheet for map
Farmland of statewide importance, if subsoiled,		not frequently flooded during the growing season	Backgrour	Local Roads	
root inhibiting soil layer Farmland of statewide		Farmland of statewide importance, if warm	Backgroui	Aerial Photography	Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)
importance, if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60		enough Farmland of statewide importance, if thawed Farmland of local importance Farmland of local importance, if irrigated			 Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required. This product is generated from the USDA-NRCS certified data as of the version date(s) listed below. Soil Survey Area: Coos County, Oregon Survey Area Data: Version 18, Sep 8, 2022 Soil map units are labeled (as space allows) for map scales 1:50,000 or larger. Date(s) aerial images were photographed: Oct 5, 2019—Oct 10, 2019 The orthophoto or other base map on which the soil lines were
					compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Farmland Classification

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
8B	Bullards sandy loam, 0 to 7 percent slopes	Farmland of statewide importance	2.5	56.0%
8E	Bullards sandy loam, 30 to 50 percent slopes	Not prime farmland	2.0	44.0%
Totals for Area of Interes	st	4.5	100.0%	

Rating Options—Farmland Classification

Aggregation Method: No Aggregation Necessary

Tie-break Rule: Lower

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Building Site Development

This folder contains a collection of tabular reports that present soil interpretations related to building site development. The reports (tables) include all selected map units and components for each map unit, limiting features and interpretive ratings. Building site development interpretations are designed to be used as tools for evaluating soil suitability and identifying soil limitations for various construction purposes. As part of the interpretation process, the rating applies to each soil in its described condition and does not consider present land use. Example interpretations can include corrosion of concrete and steel, shallow excavations, dwellings with and without basements, small commercial buildings, local roads and streets, and lawns and landscaping.

Dwellings and Small Commercial Buildings

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. This table shows the degree and kind of soil limitations that affect dwellings and small commercial buildings.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. *Not limited* indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. *Somewhat limited* indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. *Very limited* indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the table indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.01 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced

concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Information in this table is intended for land use planning, for evaluating land use alternatives, and for planning site investigations prior to design and construction. The information, however, has limitations. For example, estimates and other data generally apply only to that part of the soil between the surface and a depth of 5 to 7 feet. Because of the map scale, small areas of different soils may be included within the mapped areas of a specific soil.

The information is not site specific and does not eliminate the need for onsite investigation of the soils or for testing and analysis by personnel experienced in the design and construction of engineering works.

Government ordinances and regulations that restrict certain land uses or impose specific design criteria were not considered in preparing the information in this table. Local ordinances and regulations should be considered in planning, in site selection, and in design.

Report—Dwellings and Small Commercial Buildings

[Onsite investigation may be needed to validate the interpretations in this table and to confirm the identity of the soil on a given site. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. The table shows only the top five limitations for any given soil. The soil may have additional limitations]

Dwellings and Small Commercial Buildings–Coos County, Oregon							
Map symbol and soil	Pct. of map unit	Dwellings without bas	ements	Dwellings with base	ments	Small commercial buildings	
name		Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8B—Bullards sandy loam, 0 to 7 percent slopes							
Bullards	75	Not limited		Not limited		Somewhat limited	
						Slope	0.01
8E—Bullards sandy loam, 30 to 50 percent slopes							
Bullards	80	Very limited		Very limited		Very limited	
		Slope	1.00	Slope	1.00	Slope	1.00

Land Classifications

This folder contains a collection of tabular reports that present a variety of soil groupings. The reports (tables) include all selected map units and components for each map unit. Land classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Prime and other Important Farmlands

This table lists the map units in the survey area that are considered important farmlands. Important farmlands consist of prime farmland, unique farmland, and farmland of statewide or local importance. This list does not constitute a recommendation for a particular land use.

In an effort to identify the extent and location of important farmlands, the Natural Resources Conservation Service, in cooperation with other interested Federal, State, and local government organizations, has inventoried land that can be used for the production of the Nation's food supply.

Prime farmland is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil quality, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. The water supply is dependable and of adequate quality. Prime farmland is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

For some of the soils identified in the table as prime farmland, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

A recent trend in land use in some areas has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

Unique farmland is land other than prime farmland that is used for the production of specific high-value food and fiber crops, such as citrus, tree nuts, olives, cranberries, and other fruits and vegetables. It has the special combination of soil quality, growing season, moisture supply, temperature, humidity, air drainage, elevation, and aspect needed for the soil to economically produce sustainable high yields of these crops when properly managed. The water supply is dependable and of adequate quality. Nearness to markets is an additional consideration. Unique farmland is not based on national criteria. It commonly is in areas where there is a special microclimate, such as the wine country in California.

In some areas, land that does not meet the criteria for prime or unique farmland is considered to be *farmland of statewide importance* for the production of food, feed, fiber, forage, and oilseed crops. The criteria for defining and delineating farmland of statewide importance are determined by the appropriate State agencies. Generally, this land includes areas of soils that nearly meet the requirements for prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. Some areas may produce as high a yield as prime farmland if conditions are favorable. Farmland of statewide importance may include tracts of land that have been designated for agriculture by State law.

In some areas that are not identified as having national or statewide importance, land is considered to be *farmland of local importance* for the production of food, feed, fiber, forage, and oilseed crops. This farmland is identified by the appropriate local agencies. Farmland of local importance may include tracts of land that have been designated for agriculture by local ordinance.

Report—Prime and other Important Farmlands

Custom Soil Resource Report

	Prime and other Important Farmlands–Coos County, Oregon						
Map Symbol	Map Unit Name	Farmland Classification					
8B	Bullards sandy loam, 0 to 7 percent slopes	Farmland of statewide importance					
8E	Bullards sandy loam, 30 to 50 percent slopes	Not prime farmland					

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf



USDA United States Department of Agriculture

> Natural Resources

Conservation Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Coos County, Oregon



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map	9
Legend	10
Map Unit Legend	11
Map Unit Descriptions	11
Coos County, Oregon	13
5A—Blacklock fine sandy loam, 0 to 3 percent slopes	13
5B—Blacklock fine sandy loam, 3 to 7 percent slopes	14
8B—Bullards sandy loam, 0 to 7 percent slopes	15
8C—Bullards sandy loam, 7 to 12 percent slopes	16
8E—Bullards sandy loam, 30 to 50 percent slopes	17
28—Heceta fine sand	18
Soil Information for All Uses	20
Suitabilities and Limitations for Use	20
Land Classifications	20
Farmland Classification (Neighborhood)	20
Soil Reports	26
Land Classifications	26
Prime and other Important Farmlands (Neighborhood)	26
References	29

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



	MAP LEGEND			MAP INFORMATION
Area of Int	terest (AOI) Area of Interest (AOI)	8	Spoil Area Stony Spot	The soil surveys that comprise your AOI were mapped at 1:20,000.
Soils	Soil Map Unit Polygons Soil Map Unit Lines Soil Map Unit Points Point Features	© * -	Very Stony Spot Wet Spot Other Special Line Features	Warning: Soil Map may not be valid at this scale. Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed
() () () () () () () () () () () () () (Blowout Borrow Pit Clay Spot	Water Fea Transporta	tures Streams and Canals ation Rails	scale. Please rely on the bar scale on each map sheet for map measurements.
◇ ¥ *	Closed Depression Gravel Pit Gravelly Spot	 Intersta US Rou Major R Local R Background Aerial P 	Interstate Highways US Routes Major Roads	Source of Map: Natural Resources Conservation Service Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)
۵ بر ش	Landfill Lava Flow Marsh or swamp Mine or Quarry		Local Roads nd Aerial Photography	Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.
© 0 ~	Miscellaneous Water Perennial Water Rock Outcrop			This product is generated from the USDA-NRCS certified data as of the version date(s) listed below. Soil Survey Area: Coos County, Oregon
+ :: =	Saline Spot Sandy Spot Severely Eroded Spot			Survey Area Data: Version 18, Sep 8, 2022 Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.
۵ بو بو	Sinkhole Slide or Slip Sodic Spot			Date(s) aerial images were photographed: Oct 5, 2019—Oct 10, 2019 The orthophoto or other base map on which the soil lines were
				compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
5A	Blacklock fine sandy loam, 0 to 3 percent slopes	2.8	5.1%
5B	Blacklock fine sandy loam, 3 to 7 percent slopes	0.1	0.1%
8B	Bullards sandy loam, 0 to 7 percent slopes	22.9	41.5%
8C	Bullards sandy loam, 7 to 12 percent slopes	8.5	15.4%
8E	Bullards sandy loam, 30 to 50 percent slopes	14.8	26.9%
28	Heceta fine sand	6.1	11.0%
Totals for Area of Interest		55.1	100.0%

Map Unit Legend

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it

was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Coos County, Oregon

5A—Blacklock fine sandy loam, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 21qb Elevation: 30 to 350 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of unique importance

Map Unit Composition

Blacklock and similar soils: 75 percent Minor components: 8 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blacklock

Setting

Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Parent material: Sandy marine deposits

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

H1 - 1 to 4 inches: fine sandy loam

H2 - 4 to 16 inches: loamy fine sand

- H3 16 to 53 inches: cemented
- H4 53 to 76 inches: sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: 12 to 20 inches to ortstein
Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 0 to 18 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 2.6 inches)

Interpretive groups

Land capability classification (irrigated): 4w Land capability classification (nonirrigated): 6w Hydrologic Soil Group: C/D Ecological site: F004AC015OR - Aquic Forest Hydric soil rating: Yes

Minor Components

Blacklock, clayey substratum

Percent of map unit: 8 percent Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: Yes

5B—Blacklock fine sandy loam, 3 to 7 percent slopes

Map Unit Setting

National map unit symbol: 21qc Elevation: 0 to 350 feet Mean annual precipitation: 50 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of unique importance

Map Unit Composition

Blacklock and similar soils: 75 percent Minor components: 8 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blacklock

Setting

Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Parent material: Sandy marine deposits

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material *H1 - 1 to 4 inches:* fine sandy loam *H2 - 4 to 16 inches:* loamy fine sand *H3 - 16 to 53 inches:* cemented *H4 - 53 to 76 inches:* sand

Properties and qualities

Slope: 3 to 7 percent
Depth to restrictive feature: 12 to 20 inches to ortstein
Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 0 to 18 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 2.6 inches)

Interpretive groups

Land capability classification (irrigated): 4w Land capability classification (nonirrigated): 6w Hydrologic Soil Group: C/D *Ecological site:* F004AC015OR - Aquic Forest *Hydric soil rating:* Yes

Minor Components

Heceta

Percent of map unit: 8 percent Landform: Deflation basins on dunes Down-slope shape: Linear Across-slope shape: Linear Other vegetative classification: Poorly Drained (G004AY018OR) Hydric soil rating: Yes

8B—Bullards sandy loam, 0 to 7 percent slopes

Map Unit Setting

National map unit symbol: 21rc Elevation: 30 to 600 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Bullards and similar soils: 75 percent *Minor components:* 9 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Bullards

Setting

Landform: Marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Parent material: Mixed eolian and marine deposits

Typical profile

Oi - 0 to 3 inches: slightly decomposed plant material

H1 - 3 to 10 inches: sandy loam

H2 - 10 to 44 inches: gravelly sandy loam

H3 - 44 to 63 inches: sand

Properties and qualities

Slope: 0 to 7 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None

Frequency of ponding: None *Available water supply, 0 to 60 inches:* Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: F004AC410OR - Coastal Upland Warm Forest Forage suitability group: Well Drained <15% Slopes (G004AY014OR) Other vegetative classification: Well Drained <15% Slopes (G004AY014OR) Hydric soil rating: No

Minor Components

Blacklock

Percent of map unit: 9 percent Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: Yes

8C—Bullards sandy loam, 7 to 12 percent slopes

Map Unit Setting

National map unit symbol: 21rd Elevation: 30 to 600 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Bullards and similar soils: 75 percent *Minor components:* 8 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Bullards

Setting

Landform: Marine terraces Landform position (three-dimensional): Riser Down-slope shape: Linear Across-slope shape: Linear Parent material: Mixed eolian and marine deposits

Typical profile

Oi - 0 to 3 inches: slightly decomposed plant material *H1 - 3 to 10 inches:* sandy loam

H2 - 10 to 44 inches: gravelly sandy loam

H3 - 44 to 63 inches: sand

Properties and qualities

Slope: 7 to 12 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: F004AC410OR - Coastal Upland Warm Forest Forage suitability group: Well Drained <15% Slopes (G004AY014OR) Other vegetative classification: Well Drained <15% Slopes (G004AY014OR) Hydric soil rating: No

Minor Components

Blacklock

Percent of map unit: 8 percent Landform: Depressions on marine terraces Landform position (three-dimensional): Tread Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: Yes

8E—Bullards sandy loam, 30 to 50 percent slopes

Map Unit Setting

National map unit symbol: 21rg Elevation: 50 to 600 feet Mean annual precipitation: 55 to 75 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Not prime farmland

Map Unit Composition

Bullards and similar soils: 80 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Bullards

Setting

Landform: Marine terraces Landform position (three-dimensional): Riser Down-slope shape: Linear Across-slope shape: Linear Parent material: Mixed eolian and marine deposits

Typical profile

Oi - 0 to 3 inches: slightly decomposed plant material

H1 - 3 to 10 inches: sandy loam

H2 - 10 to 44 inches: gravelly sandy loam

H3 - 44 to 63 inches: sand

Properties and qualities

Slope: 30 to 50 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6e Hydrologic Soil Group: B Ecological site: F004AC410OR - Coastal Upland Warm Forest Hydric soil rating: No

28—Heceta fine sand

Map Unit Setting

National map unit symbol: 21n8 Elevation: 0 to 80 feet Mean annual precipitation: 50 to 70 inches Mean annual air temperature: 52 to 54 degrees F Frost-free period: 200 to 240 days Farmland classification: Farmland of statewide importance

Map Unit Composition

Heceta and similar soils: 80 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Heceta

Setting

Landform: Deflation basins on dunes Down-slope shape: Linear Across-slope shape: Linear Parent material: Eolian deposits

Typical profile

H1 - 0 to 4 inches: fine sand *H2 - 4 to 60 inches:* sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 0 inches
Frequency of flooding: None
Frequency of ponding: Frequent
Available water supply, 0 to 60 inches: Low (about 3.6 inches)

Interpretive groups

Land capability classification (irrigated): 4w Land capability classification (nonirrigated): 4w Hydrologic Soil Group: A/D Ecological site: R004AB203OR - Aquic Interdune Forage suitability group: Poorly Drained (G004AY018OR) Other vegetative classification: Poorly Drained (G004AY018OR) Hydric soil rating: Yes

Soil Information for All Uses

Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

Land Classifications

Land Classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Farmland Classification (Neighborhood)

Farmland classification identifies map units as prime farmland, farmland of statewide importance, farmland of local importance, or unique farmland. It identifies the location and extent of the soils that are best suited to food, feed, fiber, forage, and oilseed crops. NRCS policy and procedures on prime and unique farmlands are published in the "Federal Register," Vol. 43, No. 21, January 31, 1978.

Custom Soil Resource Report Map—Farmland Classification (Neighborhood)





Custom Soil Resource Report

Prime farmland if Farmland of statewide Farmland of statewide Farmland of unique Prime farmland if 1 A الريادي -----subsoiled, completely importance, if drained and importance, if irrigated importance subsoiled, completely removing the root either protected from and reclaimed of excess removing the root Not rated or not available $\mathcal{F}^{(1)}(\mathcal{F})$ inhibiting soil layer flooding or not frequently salts and sodium inhibiting soil layer flooded during the Soil Rating Points Prime farmland if irrigated Farmland of statewide Prime farmland if arowing season and the product of I (soil importance, if drained or irrigated and the product Not prime farmland erodibility) x C (climate Farmland of statewide either protected from of I (soil erodibility) x C factor) does not exceed importance, if irrigated flooding or not frequently All areas are prime (climate factor) does not and drained flooded during the farmland exceed 60 60 growing season Prime farmland if irrigated Farmland of statewide Prime farmland if drained Prime farmland if --and reclaimed of excess importance, if irrigated Farmland of statewide irrigated and reclaimed -Prime farmland if salts and sodium and either protected from importance, if warm of excess salts and protected from flooding or flooding or not frequently enough, and either sodium Farmland of statewide ----not frequently flooded flooded during the drained or either Farmland of statewide importance during the growing growing season protected from flooding or importance Farmland of statewide not frequently flooded season a 🖬 Farmland of statewide Farmland of statewide importance, if drained during the growing Prime farmland if irrigated importance, if subsoiled. importance, if drained Farmland of statewide season completely removing the importance, if protected Prime farmland if drained Farmland of statewide root inhibiting soil layer Farmland of statewide from flooding or not and either protected from importance, if protected importance, if warm Farmland of statewide 100 frequently flooded during flooding or not frequently from flooding or not enough importance, if irrigated the growing season flooded during the frequently flooded during and the product of I (soil Farmland of statewide growing season the growing season Farmland of statewide 1990 B erodibility) x C (climate importance, if thawed importance, if irrigated Prime farmland if irrigated Farmland of statewide factor) does not exceed Farmland of local 1000 and drained importance, if irrigated 60 importance Prime farmland if irrigated Farmland of local ----and either protected from importance, if irrigated flooding or not frequently flooded during the growing season

Custom Soil Resource Report

Farmland of statewide importance, if drained and either protected from flooding or not frequently		Farmland of statewide importance, if irrigated and reclaimed of excess salts and sodium		Farmland of unique importance Not rated or not available	The soil surveys that comprise your AOI were mapped at 1:20,000.							
flooded during the growing season		Farmland of statewide	Water Features		Warning: Soil Map may not be valid at this scale.							
Farmland of statewide importance, if irrigated		either protected from flooding or not frequently flooded during the growing season Farmland of statewide importance, if warm enough. and either	either protected from flooding or not frequently	either protected from flooding or not frequently	either protected from flooding or not frequently	either protected from flooding or not frequently	either protected from flooding or not frequently	either protected from flooding or not frequently	either protected from flooding or not frequently	Transporta	ation	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil
and drained Farmland of statewide	_		••••	Rails	line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed							
and either protected from flooding or not frequently			~	US Routes	scale.							
flooded during the growing season		drained or either protected from flooding or	~	Major Roads	Please rely on the bar scale on each map sheet for map							
Farmland of statewide importance, if subsoiled,		not frequently flooded during the growing season	Backgroun	Local Roads								
root inhibiting soil layer Farmland of statewide		Farmland of statewide importance, if warm	Backgroun	Aerial Photography	Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)							
importance, if irrigated and the product of I (soil erodibility) x C (climate factor) does not exceed 60		enough Farmland of statewide importance, if thawed Farmland of local importance Farmland of local importance, if irrigated			 Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required. This product is generated from the USDA-NRCS certified data as of the version date(s) listed below. Soil Survey Area: Coos County, Oregon Survey Area Data: Version 18, Sep 8, 2022 Soil map units are labeled (as space allows) for map scales 1:50,000 or larger. Date(s) aerial images were photographed: Oct 5, 2019—Oct 10, 2019 The arther base map on which the sail lines were 							
					I he orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.							

Table—Farmland Classification (Neighborhood)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
5A	Blacklock fine sandy loam, 0 to 3 percent slopes	Farmland of unique importance	2.8	5.1%
5B	Blacklock fine sandy loam, 3 to 7 percent slopes	Farmland of unique importance	0.1	0.1%
8B	Bullards sandy loam, 0 to 7 percent slopes	Farmland of statewide importance	22.9	41.5%
8C	Bullards sandy loam, 7 to 12 percent slopes	Farmland of statewide importance	8.5	15.4%
8E	Bullards sandy loam, 30 to 50 percent slopes	Not prime farmland	14.8	26.9%
28	Heceta fine sand	Farmland of statewide importance	6.1	11.0%
Totals for Area of Interes	st	55.1	100.0%	

Rating Options—Farmland Classification (Neighborhood)

Aggregation Method: No Aggregation Necessary

Tie-break Rule: Lower

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Land Classifications

This folder contains a collection of tabular reports that present a variety of soil groupings. The reports (tables) include all selected map units and components for each map unit. Land classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Prime and other Important Farmlands (Neighborhood)

This table lists the map units in the survey area that are considered important farmlands. Important farmlands consist of prime farmland, unique farmland, and farmland of statewide or local importance. This list does not constitute a recommendation for a particular land use.

In an effort to identify the extent and location of important farmlands, the Natural Resources Conservation Service, in cooperation with other interested Federal, State, and local government organizations, has inventoried land that can be used for the production of the Nation's food supply.

Prime farmland is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil quality, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. The water supply is dependable and of adequate quality. Prime farmland is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

For some of the soils identified in the table as prime farmland, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures.

A recent trend in land use in some areas has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

Unique farmland is land other than prime farmland that is used for the production of specific high-value food and fiber crops, such as citrus, tree nuts, olives, cranberries, and other fruits and vegetables. It has the special combination of soil quality, growing season, moisture supply, temperature, humidity, air drainage, elevation, and aspect needed for the soil to economically produce sustainable high yields of these crops when properly managed. The water supply is dependable and of adequate quality. Nearness to markets is an additional consideration. Unique farmland is not based on national criteria. It commonly is in areas where there is a special microclimate, such as the wine country in California.

In some areas, land that does not meet the criteria for prime or unique farmland is considered to be *farmland of statewide importance* for the production of food, feed, fiber, forage, and oilseed crops. The criteria for defining and delineating farmland of statewide importance are determined by the appropriate State agencies. Generally, this land includes areas of soils that nearly meet the requirements for prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. Some areas may produce as high a yield as prime farmland if conditions are favorable. Farmland of statewide importance may include tracts of land that have been designated for agriculture by State law.

In some areas that are not identified as having national or statewide importance, land is considered to be *farmland of local importance* for the production of food, feed, fiber, forage, and oilseed crops. This farmland is identified by the appropriate local agencies. Farmland of local importance may include tracts of land that have been designated for agriculture by local ordinance.

Prime and other Important Farmlands–Coos County, Oregon					
Map Symbol	Map Unit Name	Farmland Classification			
5A	Blacklock fine sandy loam, 0 to 3 percent slopes	Farmland of unique importance			
5B	Blacklock fine sandy loam, 3 to 7 percent slopes	Farmland of unique importance			
8B	Bullards sandy loam, 0 to 7 percent slopes	Farmland of statewide importance			
8C	Bullards sandy loam, 7 to 12 percent slopes	Farmland of statewide importance			
8E	Bullards sandy loam, 30 to 50 percent slopes	Not prime farmland			

Report—Prime and other Important Farmlands (Neighborhood)

Custom Soil Resource Report

Prime and other Important Farmlands–Coos County, Oregon			
Map Symbol	Map Unit Name	Farmland Classification	
28	Heceta fine sand	Farmland of statewide importance	

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf